

## Optical observations of meteoric dust in the middle atmosphere during Leonid activity in recent years 2001–2003 over India

B. Padma Kumari, J. M. Trigo-Rodríguez,<sup>1</sup> A. L. Londhe, D. B. Jadhav, and H. K. Trimbake

Indian Institute of Tropical Meteorology, Pune, India

Received 26 May 2005; revised 4 July 2005; accepted 25 July 2005; published 23 August 2005.

[1] Twilight photometric technique has been used in order to demonstrate that the middle atmosphere can be perturbed by the presence of tiny particles delivered after an encounter with cometary dust trails like the one produced by 55P/Tempel-Tuttle. The presence of meteoric dust in the atmosphere from the Leonid activity that occurred from 2001 to 2003 was detected by the twilight photometer operated at Pune (18.5°N, 73.9°E), India. The November 2001 and 2002 Leonid storms, and the 2003 November outburst, caused significant enhancements of dust from just above the mesopause to the lower stratosphere. The present study shows the formation of meteoric dust layers at mesospheric levels and their subsequent descent to lower altitudes. The enhanced stratospheric layers are observed 4 to 8 days after the peak meteor activity. **Citation:** Padma Kumari, B., J. M. Trigo-Rodríguez, A. L. Londhe, D. B. Jadhav, and H. K. Trimbake (2005), Optical observations of meteoric dust in the middle atmosphere during Leonid activity in recent years 2001–2003 over India, *Geophys. Res. Lett.*, 32, L16807, doi:10.1029/2005GL023434.

### 1. Introduction

[2] Meteoric dust is being continuously delivered to the terrestrial atmosphere from comets and asteroids. It is important to quantify the order of magnitude of the dust influx produced by unusual high-activity meteoric events in order to decipher its possible contribution to atmospheric and climate physical processes. The past encounters of the Earth with dust trails associated with 55P/Tempel-Tuttle presented an opportunity to study the magnitude of dust input during meteor storm and outburst conditions. Since these conditions are unusual and extreme they will provide an excellent opportunity to try and constrain the influx of debris from extraterrestrial sources and their behavior in the middle atmosphere. When the Earth intercepts a dust trail, thousands of particles ranging from tens of microns to several centimeters enter the terrestrial atmosphere at high velocity. Meteoroids from 55P/Tempel-Tuttle dust trails enter the Earth's atmosphere with an average geocentric velocity of 70.6 km/s. At this high velocity the original meteoroids are not expected to retain their original morphology, but instead survive as quenched, molten silicate spheres [Rietmeijer and Jenniskens, 2000].


Consequently, the resulting mass delivered to the atmosphere and surviving ablation is not well known. Recently, Trigo-Rodríguez *et al.* [2004a] estimated the mass influx of Leonid particles during past Leonid outbursts and storms, from the dependence of the size distribution and relative abundance of large particles. If large particles dominate the dust trail (as the 1998 outburst) the released mass would be on the order of several metric tons, whereas trails composed of fine dust particles would be one or two orders of magnitude lower depending on the intensity and extension of the high-activity level [Trigo-Rodríguez *et al.*, 2004a]. Small spheres or condensates of metal rich particles are produced during ablation of high-velocity meteoroids [Rietmeijer, 2002a, 2002b]. Another approach to this problem is the measurement of cosmic dust collected in different terrestrial environments; the recent estimate of the terrestrial mass accretion rate for meteoroids of all sizes is  $(49–56) \times 10^6$  kg/yr [Love and Brownlee, 1993]. Consequently, the Leonid input would be significant compared with the daily average for the Earth from all sources, i.e.,  $\sim 85 \times 10^3$  kg.

[3] On the rare occasions when Earth intercepts a dust trail a meteor storm is produced [Jenniskens, 1996]. Approximately every 33.3 yr the return to perihelion of comet 55P/Tempel-Tuttle makes it possible for the Earth to intercept dust trails associated with this comet. Leonid storms produce the strongest meteor rates visible at Earth [Brown and Arlt, 2000]. The recent storms associated with the last 55P/Tempel-Tuttle return to perihelion have been reasonable-well predicted [McNaught and Asher, 1999, 2001; Lyytinen and Jenniskens, 2003].

[4] On Nov. 18, 2001, the Earth crossed the 1866 dust trail, the intensity of which was reinforced by the presence of the 1669 and 1767 dust trails. So the Leonid annual shower turned into a meteor storm as had been predicted [McNaught and Asher, 1999, 2001]. A very strong shower peak of  $\sim 3400$  meteors/hr was reported in 2001 [Arlt *et al.*, 2001]. Again on Nov. 19, 2002, the Earth intercepted two 55P/Tempel-Tuttle dust trails [McNaught and Asher, 2002]. As a consequence, two strong peaks were observed in 2002 with typical spatial number densities of  $\sim 3000$  particles heavier than  $6 \times 10^{-6}$  g in a cube with 1000 km-long edges [Trigo-Rodríguez *et al.*, 2004b]. On November 2003, the Earth did not cross any dust trails, but two meteor outbursts occurred on Nov. 13 and Nov. 19 as predicted [Vaubailion *et al.*, 2003].

[5] Here we present the detection of an important input of Leonid ablation debris in the atmosphere during Nov. 2001, 2002 and 2003. We used the twilight photometric technique, a distant atmospheric sounding method used in the past for

<sup>1</sup>Now at Institute of Geophysics and Planetary Physics, University of California, Los Angeles, California, USA.



**Figure 1.** Logarithmic gradient of evening twilight intensity ( $q$ ) curves for Nov. 2001. The Leonid storm took place on Nov. 18. Nov. 16 shows the atmospheric condition before the storm. Although the profiles on 18 and 20 do not show any dust layers at high altitudes they show enhanced values compared to 16. The profiles on 21 and 22 show dust accumulation below 40 km. On 26, high altitudes still seem to be perturbed with small layer enhancements, and finally on 30 no enhancements are seen. (The threshold level of detection of photometer is  $0.001 \pm 2 \times 10^{-5}$  particles/cc at high altitudes and  $1 \pm 2 \times 10^{-2}$  particles/cc at low altitudes).

monitoring the influx of meteoric dust in the upper and middle atmosphere [Link, 1975; Matshvili *et al.*, 1999, 2000; Padma Kumari, 2004]. The twilight observations were carried out at Abastumani Astrophysical Observatory, Georgia ( $42.8^{\circ}\text{N}$ ,  $41.8^{\circ}\text{E}$ ) during Quadrantid,  $\eta$  Aquarid, Perseid, Orionid, Geminid and Leonid meteor showers. All these meteor events showed an enhancement of the atmospheric scattering ability [Matshvili *et al.*, 1999, 2000]. In the present study, a twilight photometry technique has been used for the first time from a ground station located in Pune (India) to monitor meteor dust in the middle atmosphere.

## 2. Methodology

[6] The twilight photometer consists of a simple experimental setup. It comprises a telescopic lens, a red filter peaking at 660 nm and a photomultiplier tube as a detector. The scattered light from the zenith sky is gathered from a  $1^{\circ}$  field of view. A more detailed description of the instrumentation and methodology is given elsewhere [Padma Kumari, 2004]. The dominating mechanism of formation of twilight sky brightness is light scattering in the sunlit layers of the atmosphere. The most important circumstance, which gives an altitudinal sounding ability to the twilight event, is that only a comparatively thin layer of air above the Earth's shadow contributes maximum to the sky brightness at every instant (see Rozenberg [1966] for more details). The most effective way of retrieving the aerosol layer is by the logarithmic gradient of intensity [Matshvili and Rietmeijer, 2002] i.e.,

$$q = -d(\log I)/dh$$


where  $I$  is measured twilight intensity obtained in relative units and  $h$  is the effective height of the Earth's shadow computed using the recorded time of observation from Earth's simple geometry [Padma Kumari, 2004]. The effect of Rayleigh scattering on the value of  $q$ , studied by Bigg [1956], is nearly constant and it does not give any structure on the  $q$  profile, and thus,  $I$  is assumed to be directly proportional to aerosol number density. The estimated uncertainty in the measurements should increase with altitude (as the signal gets smaller), but to avoid this, the signal level is kept constant through out the experiment by changing only the exposure area of the telescopic lens, and maintaining the output in the range 2 to 12 V. During this process no instrumental function is changed [Padma Kumari, 2004]. Hence, the error in measurement is  $\sim 1-2\%$ . The photometer-derived profiles are compared with that of lidar and the comparison is found to be very good [Padma Kumari *et al.*, 2004].

## 3. Results and Discussion

[7] The vertical profiles of the logarithmic gradient of twilight intensity ( $q$ ) obtained on 16, 18, 20, 21, 22, 26 and 30 November 2001, are shown in Figure 1. The horizontal bars in each profile represent the error in the measurements. Although the profiles on 18 and 20 do not show any enhancements at high altitudes except a narrow layer in the lower stratosphere, they show somewhat enhanced values at these high altitudes compared to the values on Nov. 16 (before Leonid storm) and Nov. 26 and 30 well after the Leonid storm. On Nov. 21, it is likely that dust particles from higher levels descended to lower ones and accumulated at altitudes below 40 km. On Nov. 22, a broad layer is observed from the tropopause to 40 km with a peak at 30 km. On Nov. 26, the higher altitudes still seem to be perturbed with small layer enhancements at about 30 km, 58 km and 70 km, indicating a further influx of meteoric dust. Finally, on Nov. 30 no enhancements are seen except for a small peak at  $\sim 20$  km.

[8] In the year 2002, the vertical profiles were obtained during the evening twilights of Nov. 18, 25, 27 and 29 (Figure 2). No significant enhancements are observed at higher altitudes in the profile obtained on Nov. 18 because the observations were performed hours before the two Leonid storms. On Nov. 25 an evident enhancement in the profile was observed from 40–55 km indicating the presence of dust layer in this altitude range. During the following days this dust layer descended to lower altitudes. In fact, the profile on Nov. 27 shows a prominent dust layer in the lower stratosphere (20–30 km). This stratospheric layer is found to decrease in intensity on Nov. 29 and, in general, no additional enhancements were observed.

[9] Despite the lower intensity of the Leonid meteor shower in 2003, the results were interesting. The vertical profiles obtained from Nov. 13 to Nov. 28, 2003 are shown in Figure 3. The profile obtained on the evening of Nov. 13 shows no obvious enhancements at high altitudes, but a stratospheric layer peaking between 20–30 km is very prominent. In the early morning of Nov. 14, the Leonid outburst peak is observed over East Asia, and on the morning of Nov. 16, a broad enhancement from 55 to 80 km peaking at  $\sim 65$  km appears in the profile. On the



**Figure 2.** Logarithmic gradient of evening twilight intensity ( $q$ ) curves for Nov. 2002. The Leonid storm took place on Nov. 19. Nov. 18 shows the atmospheric condition well before the storm. After the storm, Nov. 25 shows an enhanced layer from 40–55 km. This layer is descended to lower stratosphere by Nov. 27 and finally no enhancements are seen on Nov. 29.


morning of Nov. 17, the dust layer descended to lower altitudes, peaking between 35 and 55 km with peaks at  $\sim 40$  and  $\sim 50$  km. It seems that the layer in the profile of Nov. 16 while descending in the atmosphere broke up in two distinct dust layers. It would be produced under the influence of prevailing physical conditions in this atmospheric region. One possibility is that the dust particles were formed by a double population of different sizes (e.g. remaining ablation fragments and smaller condensates) that would be redistributed under the effect of turbulence or other physical processes. Such redistribution would be the cause of the observed differences in the intensity of both dust layers between Nov. 16 and 17. Another Leonid outburst took place on Nov. 19. This other outburst was a real independently observed phenomenon [Vaubaillon *et al.*, 2003]. As a result a broad enhancement was observed on the morning of Nov. 20 between 65 and 95 km, peaking at  $\sim 80$  km. On the morning of Nov. 25, another enhancement was observed from 20 to 50 km. Finally, on the evening of Nov. 28 no additional enhancements were seen at high altitudes, except for a narrow layer at  $\sim 20$  km. The pioneering work by Mateshvili *et al.* [2000] during the 1998 and 1999 Leonid meteor showers showed the formation of meteoric dust layers in addition to an overall enhancement of atmospheric turbidity by using the same technique.

[10] During atmospheric deceleration, Leonid meteoroids are ablated in the upper atmosphere. As a consequence of ablation, a column of ionized elements produces the luminous phenomenon known as meteors. Typical beginning and ending heights for visual Leonid meteors are respectively 120 and 90 km [Trigo-Rodríguez *et al.*, 2002]. Small meteoroid fragments survive ablation [Borovička and Jenniskens, 2000]. Previously, Mateshvili *et al.* [2000] estimated the particle diameters to range from 0.02 to  $\sim 22 \mu\text{m}$ . The presence of layers at different altitudes and their evolution as a function of time supports the idea that ablation of Leonid meteoroids produces particles in a wide

range of sizes as previously observed by Mateshvili *et al.* [1999, 2000].

[11] The appearance of different layers suggests that large objects that probably formed from incompletely melted materials several tens of microns in size were released. As a consequence of compaction during heating, ablation can create dense particles. Even fluffy, low-velocity meteoroids may melt and become dense spheres [Love and Brownlee, 1990]. The dense particles may form as condensates of meteor vapor, appearing as dust layer at higher altitudes, which can be detected as an enhancement in the twilight vertical profile. This dust layer slowly descends to lower altitudes and disappear within a few days as seen in Figures 1, 2, and 3. Probably, simple sedimentation alone cannot account for the observed rapid clearing of the upper atmosphere, but dynamics and poorly understood processes such as coagulation or electrostatic charging of the particles might also play a significant role.

[12] Finally, it is interesting to note that the detection of the meteoritic contribution was made possible by the present quiescent condition of the middle atmosphere. For example, such detection would not have been possible after the volcano Mt. Pinatubo erupted in the Philippines in 1991, which produced a large impact at stratospheric altitudes. The stratospheric aerosol depth following the Pinatubo eruption increased by about 2 orders of magnitude, making difficult to study the meteoritic contribution to the atmospheric aerosol. Since then, there have been no major volcanic eruptions. The stratosphere is presently in a quiescent condition. Satellite observations during the volcanically quiescent period have shown an increase in the background stratospheric aerosol layer [Thomason *et al.*,



**Figure 3.** Logarithmic gradient of morning (M) and evening (E) twilight intensity ( $q$ ) curves for November 2003. The Leonid outburst took place on the early morning of Nov. 14 over East Asia. Nov. 13 shows the atmospheric condition well before the outburst. After the outburst enhanced layer is seen on 16 morning and is descended by  $\sim 20$  km on 17 morning. Again a broad enhancement seen between 65–95 km on 20 morning is due to another Leonid outburst on 19. This layer is descended to lower altitudes by 25 Nov. and finally no enhancements are seen on 28 Nov.



1997]. But, there has been considerable uncertainty as to whether this increase is caused by anthropogenic activity or if the trend is influenced by volcanic eruptions. From our data it is evident that the meteoric dust particles, after high meteor activity, may contribute significantly to the physico-chemical processes taking place at stratospheric altitudes.

#### 4. Conclusions

[13] The twilight photometric technique is a useful tool for monitoring the influx of meteoric dust qualitatively in the middle atmosphere and its subsequent descent to lower altitudes. Our data show that the higher altitudes were highly perturbed after the Leonid meteor activity of 2001, 2002 and 2003. In general, 4 to 8 days after the peak meteor activity, an enhanced stratospheric layer was detected between 20 and 30 km and, sometimes, between 20 and 40 km. This layer seems to be a short-lived feature probably caused by rapid settling of ablation products of Leonid meteoroids. A couple of weeks after the meteor activity, the atmosphere recovers its normal dust distribution profile although a dust layer at 30 km altitude can persist for more time. Consequently, under the present or future quiescent periods, study of the meteoric contribution to the atmospheric aerosol layers makes sense in order to improve our knowledge of its influence on atmospheric chemistry, dynamics, and reflectivity changes.

[14] **Acknowledgment.** The authors thank Dr. P. C. S. Devara and Dr. G. B. Pant, Director IITM, for their encouragement in carrying out this work and also Department of Science and Technology (Grant No. ESS/63/155/98) for providing funds for the development of twilight photometer.

#### References

- Arlt, R., J. Kac, V. Krumov, A. Buchmann, and J. Verbert (2001), Bulletin 17 of the international Leonid watch: First global analysis of the 2001 Leonid storms, *WGN J. Int. Meteor. Organ.*, *29*, 187–194.
- Bigg, E. K. (1956), Detection of atmospheric dust and temperature inversions by twilight Scattering, *J. Meteorol.*, *13*, 262–268.
- Borovička, J., and P. Jenniskens (2000), Time resolved spectroscopy of a Leonid fireball afterglow, *Earth Moon Planets*, *82–83*, 399–428.
- Brown, P., and R. Arlt (2000), Detailed visual observations and modeling of the 1998 Leonid shower, *Mon. Not. R. Astron. Soc.*, *319*, 419–428.
- Jenniskens, P. (1996), Meteor stream activity: III. Measurement of the first in a new series of Leonid outbursts, *Meteorit. Planet. Sci.*, *31*, 177–184.
- Link, F. (1975), In the presence of cosmic dust in the upper atmosphere, *Planet. Space Sci.*, *23*, 1011–1012.
- Love, S. G., and D. Brownlee (1990), Heating and thermal transformation of micrometeoroids entering the Earth's atmosphere, *Icarus*, *89*, 26–43.
- Love, S. G., and D. Brownlee (1993), A direct measurement of the terrestrial mass accretion rate of cosmic dust, *Science*, *262*, 550–553.
- Lyytinen, E., and P. Jenniskens (2003), Meteor outbursts from long-period comet dust trails, *Icarus*, *162*, 443–452.
- Matashvili, N., G. G. Matashvili, I. Matashvili, L. Gheondjian, and O. Avsajanishvili (1999), Vertical distribution of dust particles in the Earth's atmosphere during the 1998 Leonids, *Meteorit. Planet. Sci.*, *34*, 969–973.
- Matashvili, N., G. G. Matashvili, I. Matashvili, L. Gheondjian, and Z. Kapanadze (2000), Dust particles in the atmosphere during the Leonid meteor showers of 1998 and 1999, *Earth Moon Planets*, *82–83*, 489–504.
- Matashvili, N., and F. J. M. Rietmeijer (2002), Stratospheric dust loading from early 1981 to September 1985 based on the twilight sounding method and stratospheric dust collections, *J. Volcanol. Geotherm. Res.*, *120*, 55–69.
- McNaught, R., and D. J. Asher (1999), Leonid dust trails and meteor storms, *WGN J. Int. Meteor. Organ.*, *27*, 85–102.
- McNaught, R., and D. J. Asher (2001), The 2001 Leonids and dust trail radiants, *WGN J. Int. Meteor. Organ.*, *29*, 156–164.
- McNaught, R., and D. J. Asher (2002), Leonid dust trail structure and predictions for 2002, *WGN J. Int. Meteor. Organ.*, *30*, 132–143.
- Padma Kumari, B. (2004), Study of stratospheric aerosol by passive remote sensing technique, Ph.D thesis, Pune Univ., Pune, India.
- Padma Kumari, B., A. L. Londhe, H. K. Trimbake, and D. B. Jadhav (2004), Comparison of aerosol vertical profiles derived by passive and active remote sensing techniques—A case study, *J. Atmos. Environ.*, *38*, 6679–6685.
- Rietmeijer, F. J. M. (2002a), Shower meteoroids: Constraints from interplanetary dust particles and Leonid meteors, *Earth Moon Planets*, *88*, 35–58.
- Rietmeijer, F. J. M. (2002b), The earliest chemical dust evolution in the solar nebula, *Chemie Erde*, *62*(1), 1–45.
- Rietmeijer, F. J. M., and P. Jenniskens (2000), Recognizing Leonid meteoroids among the collected stratospheric dust, *Earth Moon Planets*, *82–83*, 505–524.
- Rozenberg, G. V. (1966), *Twilight*, 380 pp., Springer, New York.
- Thomason, L. W., G. S. Kent, C. R. Trepte, and L. R. Poole (1997), A comparison of the stratospheric aerosol background periods of 1979 and 1989–1991, *J. Geophys. Res.*, *102*, 3611–3616.
- Trigo-Rodríguez, J. M., J. Llorca, and J. Fabregat (2002), On the origin of the 1999 Leonid storm as deduced from photographic observations, *Earth Moon Planets*, *91*, 107–119.
- Trigo-Rodríguez, J. M., J. Llorca, and J. Oró (2004a), Chemical abundances of cometary meteoroids from meteor spectroscopy: Implications to the Earth enrichment, in *Life in the Universe: From the Miller Experiment to the Search for Life in Other Worlds*, edited by J. Seckbach et al., pp. 201–204, Springer, New York.
- Trigo-Rodríguez, J. M., J. Llorca, E. Lyytinen, J. L. Ortiz, A. Sánchez Caso, C. Pineda, and S. Torrell (2004b), 2002 Leonid storm fluxes and related orbital elements, *Icarus*, *171*, 219–228.
- Vaubailion, J., E. Lyytinen, M. Nissinen, and D. J. Asher (2003), The 2003 Leonid shower from different approaches, *WGN J. Int. Meteor. Organ.*, *31*, 131–134.
- D. B. Jadhav, A. L. Londhe, B. Padma Kumari, and H. K. Trimbake, Indian Institute of Tropical Meteorology, Pune, 411 008, India. (padma@tropmet.res.in)
- J. M. Trigo-Rodríguez, Institute of Geophysics and Planetary Physics, University of California, Los Angeles, CA 90095-1567, USA.